

Eating well with fatty liver

Metabolic dysfunction-associated steatotic liver disease (MASLD)

Mediterranean-style eating is the recommended approach. If you are overweight, losing weight slowly can reduce liver fat.

Half:
non-starchy
vegetables



A quarter:
lean protein

A quarter:
whole grains, fruit
or starchy vegetables

Foods to enjoy



Non-starchy vegetables

At least 5 servings of vegetables and fruit a day



Lean protein



Fruit



Olive oil



Whole grains



Nuts and seeds



Fatty fish

Omega-3 fats



Avocado

Foods to limit



Sweet drinks

Soda, sports drinks, sweet tea, juice



Added sugar

Under 9 tsp a day for men, 6 tsp for women



Alcohol

Experts advise not drinking at all



Fried food



Processed sweets

Often contain trans fat



High-fat dairy

Choose low-fat or nonfat



White bread and rice

Choose whole grains



Potatoes

High-glycemic, keep portions small

If you are losing weight

- Aim for ½ to 2 pounds a week. Avoid crash diets and fasting.
- Losing 5 to 10% of your body weight can improve your liver.
- Keto and paleo diets are not proven for long-term weight loss.

If you have cirrhosis

- Your food needs are different. Ask us before you change your diet or try to lose weight.

General information for adults, not a personal diet plan. Follow the advice your own care team gave you, and talk to them before making big changes.

Questions? Call (718) 615-0400.

Medical emergency? Call 911.

Dr. K. Iswara,
Gastroenterology,
Calko Medical Center

Sources

- American Gastroenterological Association, MASLD and lifestyle changes

- American Association for the Study of Liver Diseases, Practice Guidance, 2023
- National Institute of Diabetes and Digestive and Kidney Diseases